

Classification of Supersymmetries

Victor G. Kac*

Abstract

In the first part of my talk I will explain a solution to the extension of Lie's problem on classification of "local continuous transformation groups of a finite-dimensional manifold" to the case of supermanifolds. (More precisely, the problem is to classify simple linearly compact Lie superalgebras, i.e. topological Lie superalgebras whose underlying space is a topological product of finite-dimensional vector spaces). In the second part I will explain how this result is used in a classification of superconformal algebras. The list consists of affine superalgebras and certain super extensions of the Virasoro algebra. In the third part I will discuss representation theory of affine superalgebras and its relation to "almost" modular forms. Furthermore, I will explain how the quantum reduction of these representations leads to a unified representation theory of super extensions of the Virasoro algebra. In the forth part I will discuss representation theory of exceptional simple infinite-dimensional linearly compact Lie superalgebras and will speculate on its relation to the Standard Model.

Introduction

The theory of Lie groups and Lie algebras began with the 1880 paper [L] of S. Lie where he posed the problem of classification of "local continuous transformation groups of a finite-dimensional manifold" M and gave a solution to this problem when $\dim M = 1$ and 2 .

The most important part of Lie's problem is the classification of the corresponding Lie algebras of vector fields on M up to "formal" isomorphism. A more invariant (independent of M) formulation is to classify linearly compact Lie algebras, i.e., topological Lie algebras whose underlying space is a topological product of discretely topologized finite-dimensional vector spaces [GS], [G2]. (Of course, it is well-known that it is impossible to classify even all finite-dimensional Lie algebras. What is usually meant by a "classification" is a complete list of simple algebras (no non-trivial ideals) and a description of semisimple algebras (no abelian ideals) in terms of simple ones.)

*Department of Mathematics, MIT, Cambridge, MA 02139, USA. E-mail: kac@math.mit.edu

It turned out that a solution to this problem requires quite different methods in the cases of finite-dimensional and infinite-dimensional groups. The most important advance in the finite-dimensional case was made by W. Killing and E. Cartan at the end of the 19th century who gave the celebrated classification of simple finite-dimensional Lie algebras over $\mathbb{C}$. The infinite-dimensional case was studied by Cartan in a series of papers written in the beginning of the 20th century, which culminated in his classification of infinite-dimensional “primitive” Lie algebras of vector fields on a finite-dimensional manifold [C].

The advent of supersymmetry in theoretical physics in the 1970s motivated work on the “super” extension of Lie’s problem. In the finite-dimensional case the latter problem was settled in [K2]. However, it took another 20 years before the problem was solved in the infinite-dimensional case [K7], [CK2], [CK3].

In the first part of my talk I will explain the classification of simple linearly compact Lie superalgebras. Remarkably, unlike in the Lie algebra case, the approach, based on the ideas of the papers [GS], [W], [K1] and [G2], is very similar in the finite- and infinite-dimensional cases.

The advent of conformal field theory in the mid-1980s motivated the work on classification and representation theory of superconformal algebras. In the second part of my talk I will explain how the classification of infinite-dimensional simple linearly compact Lie algebras is applied to classification of “linear” simple superconformal algebras. A complete list consists of the affine superalgebras, and of several series and one exceptional example of super extensions of the Virasoro algebra [FK]. (The most famous of these super extensions is the $N = 2$ superconformal algebra, which plays a fundamental role in the mirror symmetry theory.)

In the third part of my talk I will discuss representation theory of affine superalgebras [KW3], [KW4]. The key property of “admissible” representations of affine algebras is that their characters are modular functions. This is not so in the super case—for some mysterious reason, modular functions get replaced by closely related but more general functions, like Appell’s function [KW4].

Next, I will explain how the quantum reduction of “admissible” representations of affine superalgebras leads to a unified representation theory of (not necessarily linear) super extensions of the Virasoro algebra [KRW], [KW5]. This gives rise to a large class of supersymmetric rational conformal field theories.

In the last part of my talk I will discuss representation theory of exceptional infinite-dimensional simple linearly compact Lie superalgebras [KR1]–[KR4]. I am convinced that this theory may have applications to “real” physics. The main reason for this belief is the exceptional Lie superalgebra $E(3|6)$ whose maximal compact group of automorphisms is the gauge group of the Standard Model (= a quotient of $SU_3 \times SU_2 \times U_1$ by a cyclic group of order 6). Furthermore, representation theory of $E(3|6)$ accurately predicts the number of generations of leptons (= 3), but not so accurately the number of generations of quarks (= 5) [KR2]. It is also striking that the inclusion of the gauge group of the Standard Model in SU_5 , which is the gauge group of the Grand Unified Model, extends to the inclusion of $E(3|6)$ in $E(5|10)$, the largest exceptional linearly compact Lie superalgebra.

1. Classification of simple linearly compact Lie superalgebras.

1.1. First, recall some basic superalgebra terminology. A *superalgebra* is simply a $\mathbb{Z}/2\mathbb{Z}$ -graded algebra:

$$S = S_{\bar{0}} + S_{\bar{1}}, \text{ where } S_{\alpha}S_{\beta} \subset S_{\alpha+\beta}, \alpha, \beta \in \mathbb{Z}/2\mathbb{Z} = \{\bar{0}, \bar{1}\}.$$

If $a \in S_{\alpha}$, one says that the *parity* $p(a)$ is equal to α . A *derivation* D of parity $p(D)$ of a superalgebra S is a vector space endomorphism satisfying condition

$$D(ab) = (Da)b + (-1)^{p(D)p(a)}a(Db).$$

The sum $\text{Der } S$ of the spaces of derivations of parity $\bar{0}$ and $\bar{1}$ is closed under the super bracket:

$$[D, D_1] = DD_1 - (-1)^{p(D)p(D_1)}D_1D.$$

This super bracket satisfies super analogs of anticommutativity and Jacobi identity, hence defines what is called a *Lie superalgebra*. (The super anticommutativity axiom is $[a, b] = -(-1)^{p(a)p(b)}[b, a]$, and the super Jacobi identity axiom means that the operator $(\text{ad } a)b := [a, b]$ is a derivation.)

One of the basic constructions is the *superization* which basically amounts to adding anticommuting indeterminates. In other words, given an algebra (associative or Lie) $\mathcal{A}$ we consider the Grassmann algebra $\mathcal{A}\langle n \rangle$ in n anticommuting indeterminates $\xi_1, \dots, \xi_n$ over $\mathcal{A}$. This algebra carries a canonical $\mathbb{Z}/2\mathbb{Z}$ -gradation defined by letting $p(\mathcal{A}) = \bar{0}$, $p(\xi_i) = \bar{1}$. If $\mathcal{O}_m$ denotes the algebra of formal power series over $\mathbb{C}$ in m commuting indeterminates $x = (x_1, \dots, x_m)$ and n anticommuting indeterminates $\xi = (\xi_1, \dots, \xi_n)$: $x_i x_j = x_j x_i$, $x_i \xi_j = \xi_j x_i$, $\xi_i \xi_j = -\xi_j \xi_i$.

Note that the associative superalgebra $\mathcal{O}_m\langle n \rangle$ is linearly compact with respect to the topology for which the powers of the augmentation ideal $(x_1, \dots, x_m, \xi_1, \dots, \xi_n)$ form a fundamental system of neighborhoods of 0. The algebra $\mathcal{A}\langle n \rangle$ has odd (i.e., of parity $\bar{1}$) derivations $\partial/\partial\xi_i$ defined by

$$\frac{\partial}{\partial\xi_i}(a) = 0 \text{ for } a \in \mathcal{A}, \quad \frac{\partial}{\partial\xi_i}(\xi_j) = \delta_{ij},$$

and these derivations anticommute, i.e., $[\partial/\partial\xi_i, \partial/\partial\xi_j] = 0$.

The first basic example of a linearly compact Lie superalgebra is the Lie superalgebra denoted by $W(m|n)$, of all continuous derivations of the topological superalgebra $\mathcal{O}_m\langle n \rangle$:

$$W(m|n) = \left\{ \sum_{i=1}^m P_i(x, \xi) \frac{\partial}{\partial x_i} + \sum_{j=1}^n Q_j(x, \xi) \frac{\partial}{\partial \xi_j} \right\},$$

where $P_i(x, \xi), Q_j(x, \xi) \in \mathcal{O}_m\langle n \rangle$. In a more geometric language, this is the Lie superalgebra of all formal vector fields on a supermanifold of dimension $(m|n)$.

1.2. Cartan's theorem [C] states that a complete list of infinite-dimensional linearly compact simple Lie algebras over $\mathbb{C}$ consists of four series: the Lie algebra $W_m (= W(m|0))$ of all formal vector fields on an m -dimensional manifold, and its subalgebras S_m of divergenceless vector fields ($m > 1$), H_m of Hamiltonian vector fields (m even), K_m of contact vector fields (m odd).

There is a unique way to extend *divergence* from W_m to $W(m|n)$ such that the divergenceless vector fields form a subalgebra:

$$\operatorname{div} \left(\sum_i P_i \frac{\partial}{\partial x_i} + \sum_j Q_j \frac{\partial}{\partial \xi_j} \right) = \sum_i \frac{\partial P_i}{\partial x_i} + \sum_j (-1)^{p(Q_j)} \frac{\partial Q_j}{\partial \xi_j},$$

and the super analog of S_m is

$$S(m|n) = \{X \in W(m|n) \mid \operatorname{div} X = 0\}.$$

In order to define super analogs of the Hamiltonian and contact Lie algebras H_m and K_m , introduce a super analog of the algebra of differential forms [K2]. This is an associative superalgebra over $\mathcal{O}_m \langle n \rangle$, denoted by $\Omega(m|n)$, on generators $dx_1, \dots, dx_m, d\xi_1, \dots, d\xi_n$ and defining relations: $dx_i dx_j = -dx_j dx_i$, $dx_i d\xi_j = d\xi_j dx_i$, $d\xi_i d\xi_j = d\xi_j d\xi_i$, and the $\mathbb{Z}/2\mathbb{Z}$ -gradation defined by: $p(x_i) = p(d\xi_j) = \bar{0}$, $p(\xi_j) = p(dx_i) = \bar{1}$. This superalgebra is linearly compact in the topology defined by powers of the augmentation ideal. The topological superalgebra $\Omega(m|n)$ carries a unique continuous derivation d of parity $\bar{1}$ such that $d(x_i) = dx_i$, $d(\xi_j) = d\xi_j$, $d(dx_i) = 0$, $d(d\xi_j) = 0$. The operator d has all the usual properties, e.g.: $df = \sum_i dx_i \frac{\partial f}{\partial x_i} + \sum_j d\xi_j \frac{\partial f}{\partial \xi_j}$ for $f \in \mathcal{O}_m \langle n \rangle$, and $d^2 = 0$. As usual, for any $X \in W(m|n)$ one defines a derivation ι_X (contraction along X) of the superalgebra $\Omega(m|n)$ by the properties (here x stands for x and ξ): $p(\iota_X) = p(X) + \bar{1}$, $\iota_X(x_j) = 0$, $\iota_X(dx_j) = (-1)^{p(X)} X(x_j)$. The action of any $X \in W(m|n)$ on $\mathcal{O}_m \langle n \rangle$ extends in a unique way to the action by a derivation of $\Omega(m|n)$ such that $[X, d] = 0$. This is called Lie's derivative and is usually denoted by L_X , but we shall write X in place of L_X unless confusion may arise. One has the usual Cartan's formula for this action: $L_X = [d, \iota_X]$.

Using this action, one can define super-analogs of the Hamiltonian and contact Lie algebras for any $n \in \mathbb{Z}_+$:

$$H(m|n) = \{X \in W(m|n) \mid X\omega_s = 0\},$$

where $m = 2k$ and $\omega_s = \sum_{i=1}^k dx_i \wedge dx_{k+i} + \sum_{j=1}^n (d\xi_j)^2$,

$$K(m|n) = \{X \in W(m|n) \mid X\omega_c = f\omega_c\},$$

where $m = 2k + 1$, $\omega_c = dx_m + \sum_{i=1}^k x_i dx_{k+i} + \sum_{j=1}^n \xi_j d\xi_j$, and $f \in \mathcal{O}_m \langle n \rangle$.

Note that $W(0|n)$, $S(0|n)$ and $H(0|n)$ are finite-dimensional Lie superalgebras. The Lie superalgebras $W(0|n)$ and $S(0|n)$ are simple iff $n \geq 2$ and $n \geq 3$, respectively. However, $H(0|n)$ is not simple as its derived algebra $H'(0|n)$ has codimension 1 in $H(0|n)$, but $H'(0|n)$ is simple iff $n \geq 4$. Thus, in the Lie superalgebra

case the lists of simple finite- and infinite-dimensional algebras are much closer related than in the Lie algebra case.

These four series of Lie superalgebras are infinite-dimensional if $m \geq 1$, in which case they are simple except for $S(1|n)$. The derived algebra $S'(1|n)$ has codimension 1 in $S(1|n)$, and $S'(1|n)$ is simple iff $n \geq 2$.

Remarkably it turned out that the above four series do not exhaust all infinite-dimensional simple linearly compact Lie superalgebras (as has been suggested in [K2]). Far from it!

As was pointed out by several mathematicians, the Schouten bracket [SV] makes the space of polyvector fields on a m -dimensional manifold into a Lie superalgebra. The formal analog of this is the following fifth series of superalgebras, called by physicists the Batalin-Vilkoviski algebra (H stands here for “Hamiltonian” and O for “odd”):

$$HO(m|m) = \{X \in W(m|m) | X\omega_{os} = 0\},$$

where $\omega_{os} = \sum_{i=1}^m dx_i d\xi_i$ is an “odd” symplectic form. Furthermore, unlike in the $H(m|n)$ case, not all vector fields of $HO(m|m)$ have zero divergence, which gives rise to the sixth series:

$$SHO(m|m) = \{X \in HO(m|m) | \operatorname{div} X = 0\}.$$

The seventh series is the odd analog of $K(m|n)$ [ALS]:

$$KO(m|m+1) = \{X \in W(m|m+1) | X\omega_{oc} = f\omega_{oc}\},$$

where $\omega_{oc} = d\xi_{m+1} + \sum_{i=1}^m (\xi_i dx_i + x_i d\xi_i)$ is an odd contact form. One can take again the divergence 0 vector fields in $KO(m|m+1)$ in order to construct the eighth series, but the situation is more interesting. It turns out that for each $\beta \in \mathbb{C}$ one can define the deformed divergence $\operatorname{div}_\beta X$ [Ko], [K7], so that $\operatorname{div} = \operatorname{div}_0$ and

$$SKO(m|m+1; \beta) = \{X \in KO(m|m+1) | \operatorname{div}_\beta X = 0\}$$

is a subalgebra. The superalgebras $HO(m|m)$ and $KO(m|m+1)$ are simple iff $m \geq 2$ and $m \geq 1$, respectively. The derived algebra $SHO'(m|m)$ has codimension 1 in $SHO(m|m)$, and it is simple iff $m \geq 3$. The derived algebra $SKO'(m|m+1; \beta)$ is simple iff $m \geq 2$, and it coincides with $SKO(m|m+1; \beta)$ unless $\beta = 1$ or $\frac{m-2}{m}$ when it has codimension 1.

Some of the examples described above have simple “filtered deformations”, all of which can be obtained by the following simple construction. Let L be a subalgebra of $W(m|n)$, where n is even. Then it happens in three cases that

$$L^\sim := (1 + \Pi_{j=1}^n \xi_j) L$$

is different from L , but is closed under bracket. As a result we get the following three series of superalgebras: $S^\sim(0|n)$ [K2], $SHO^\sim(m|m)$ [CK2] and $SKO^\sim(m|m+1; \frac{m+2}{m})$ [Ko] (the constructions in [Ko] and [CK2] were more complicated). We thus get the ninth and the tenth series of simple infinite-dimensional Lie superalgebras:

$$\begin{aligned} SHO^\sim(m|m), \quad m \geq 2, m \text{ even}, \\ SKO^\sim(m|m+1; 1+2/m), \quad m \geq 3, m \text{ odd}. \end{aligned}$$

A surprising discovery was made in [Sh1] where the existence of three exceptional simple infinite-dimensional Lie superalgebras was announced. The proof of the existence along with one more exceptional example was given in [Sh2]. An explicit construction of these four examples was given later in [CK3]. The fifth exceptional example was found in the work on conformal algebras [CK1] and independently in [Sh2]. (The alleged sixth exceptional example $E(2|2)$ of [K7] turned out to be isomorphic to $SKO(2|3; 1)$ [CK3].)

Now I can state the first main theorem.

Theorem 1. [K7] *The complete list of simple infinite-dimensional linearly compact Lie superalgebras consists of ten series of examples described above and five exceptional examples: $E(1|6)$, $E(3|6)$, $E(3|8)$, $E(4|4)$, and $E(5|10)$.*

Here and before the notation $X(m|n)$ means that this superalgebra can be embedded in $W(m|n)$ and that this embedding is minimal possible; E stands for “exceptional”.

Remark. The local classification of transitive primitive (i.e., leaving no invariant fibrations) actions on a (super)manifold M is equivalent to the classification of all “primitive” pairs (L, L_0) , where L is a linearly compact Lie (super)algebra and L_0 is a maximal open subalgebra (such that $\dim L/L_0 = \dim M$) without non-zero ideals of L . If L is simple, choosing any maximal open subalgebra L_0 , we get a primitive pair (L, L_0) . One can show that if, in addition, L is a Lie algebra and $\dim L = \infty$, there exists a unique such L_0 . According to Cartan’s theorem, the remaining infinite-dimensional primitive pairs are the Lie algebras obtained from S_n and H_n by adding the Euler operator E . Using the structure results on general transitive linearly compact Lie algebras [G1], it is not difficult to reduce the classification of infinite-dimensional primitive pairs to the classification of simple infinite-dimensional linearly compact Lie algebras (cf. [G2]). Such a reduction is possible also in the Lie superalgebra case, but it is much more complicated for two reasons: (a) a simple linearly compact Lie superalgebra may have several maximal open subalgebras (see [CK3] for a classification), (b) construction of arbitrary primitive pairs in terms of simple primitive pairs is more complicated in the superalgebra case (see [K8]).

1.3. Here I will describe the classification of finite-dimensional simple Lie superalgebras. We already have four “non-classical” series: $W(0|n)$, $S(0|n)$, $H'(0|n)$ and $S^\sim(0|n)$. The four “classical” series are constructed as follows. Introduce the following even and odd Euler operators $E = \sum_i x_i \frac{\partial}{\partial x_i} + \sum_j \xi_j \frac{\partial}{\partial \xi_j} \in W(m|n)$ and $E_o = \sum_i x_i \frac{\partial}{\partial \xi_i} + \sum_i \xi_i \frac{\partial}{\partial x_i} \in W(m|m)$. Let

$$\begin{aligned} s\ell(m|n) &= \{X \in S(m|n) | [E, X] = 0\}, \\ spo(m|n) &= \{X \in H(m|n) | [E, X] = 0\}, \\ p(m|m) &= \{X \in SHO(m|m) | [E, X] = 0\}, \\ q(m|m) &= \{X \in W(m|m) | [E_o, X] = 0\}. \end{aligned}$$

The Lie algebras $sl_m = sl(m|0)$, $sp_m = spo(m|0)$, $so_n = spo(0|n)$ are simple. Furthermore, $sl(m|n)$ are simple for $m \neq n$, all $spo(m|n)$ and $p(m|m)$ ($m \geq 3$) are simple. The superalgebra $sl(m|m)$ contains 1-dimensional ideal $\mathbb{C}E$ and $sl(m|m)/\mathbb{C}E$ is simple for $m \geq 2$. Finally, the derived algebra $q'(m|m)$ has codimension 1 in $q(m|m)$ and $q'(m|m)/\mathbb{C}E$ is simple for $m \geq 3$.

Theorem 2. [K2] *The complete list of simple finite-dimensional Lie superalgebras consists of eight series of examples described above, the exceptional Lie superalgebras $F(4)$ and $G(3)$ of dimension 40 and 31, respectively, a 1-parameter family of 17-dimensional exceptional Lie superalgebras $D(2, 1; a)$, and the five exceptional Lie algebras.*

1.4. Plan of the proof of Theorem 1.

Step 1. Introduce Weisfeiler's filtration [W] of a simple linearly compact Lie superalgebra L . For that choose a maximal open subalgebra L_0 of L and a minimal subspace L_{-1} satisfying the properties: $L_{-1} \supsetneq L_0$, $[L_0, L_{-1}] \subset L_{-1}$. (Geometrically this corresponds to a choice of a primitive action of L and an invariant irreducible differential system.) The pair L_{-1}, L_0 can be included in a unique filtration: $L = L_{-d} \supsetneq L_{-d+1} \supset \cdots \supset L_{-1} \supset L_0 \supset L_1 \supset \cdots$, called *Weisfeiler's filtration* of *depth* d . (In the Lie algebra case, $d > 1$ only for K_m , when $d = 2$, but in the Lie superalgebra case, $d > 1$ in the majority of cases.) The associated to Weisfeiler's filtration $\mathbb{Z}$ -graded Lie superalgebra is of the form $GrL = \Pi_{j \geq -d} \mathfrak{g}_j$, and has the following properties:

- (G0) $\dim \mathfrak{g}_j < \infty$ (since $\text{codim } L_0 < \infty$),
- (G1) $\mathfrak{g}_{-j} = \mathfrak{g}_{-1}^j$ for $j \geq 1$ (by maximality of L_0),
- (G2) $[x, \mathfrak{g}_{-1}] = 0$ for $x \in \mathfrak{g}_j$, $j \geq 0 \Rightarrow x = 0$ (by simplicity of L),
- (G3) $\mathfrak{g}_0$ -module $\mathfrak{g}_{-1}$ is irreducible (by choice of L_{-1} , and faithful (by (G2)).

Weisfeiler's idea was that property (G3) is so restrictive, that it should lead to a complete classification of $\mathbb{Z}$ -graded Lie algebras satisfying (G0)–(G3). (Incidentally, the infinite-dimensionality of L and hence of GrL , since L is simple, is needed only in order to conclude that $\mathfrak{g}_1 \neq 0$.) This indeed turned out to be the case [K1]. In fact, my idea was to replace the condition of finiteness of the depth by finiteness of the growth, which allowed one to add to the Lie-Cartan list some new Lie algebras, called nowadays affine Kac-Moody algebras.

However, unlike in the Lie algebra case, it is impossible to classify all finite-dimensional irreducible faithful representations of Lie superalgebras. One needed a new idea to make this approach work.

Step 2. The main new idea is to choose L_0 to be invariant with respect to all inner automorphisms of L (meaning to contain all even ad-exponentiable elements of L). A non-trivial point is the existence of such L_0 . This is proved by making use of the characteristic supervariety, which involves rather difficult arguments of Guillemin [G2].

Next, using a normalizer trick of Guillemin [G2], I prove, for this choice of L_0 , the following very powerful restriction on the $\mathfrak{g}_0$ -module $\mathfrak{g}_{-1}$ (at this point $\dim L = \infty$ is used):

(G4) $[\mathfrak{g}_0, x] = \mathfrak{g}_{-1}$ for any non-zero even element x of $\mathfrak{g}_{-1}$.

Step 3. Consider a faithful irreducible representation of a Lie superalgebra $\mathfrak{p}$ in a finite-dimensional vector space V . This representation is called *strongly transitive* if $\mathfrak{p} \cdot x = V$ for any non-zero even element $x \in V$. By properties (G0), (G3) and (G4), the $\mathfrak{g}_0$ -module $\mathfrak{g}_{-1}$ is strongly transitive.

In order to demonstrate the power of this restriction, consider first the case when $\mathfrak{p}$ is a Lie algebra and V is purely even. Then the strong transitivity simply means that $V \setminus \{0\}$ is a single orbit of the Lie group P corresponding to $\mathfrak{p}$. It is rather easy to see that the only strongly transitive subalgebras $\mathfrak{p}$ of $\mathfrak{gl}_V$ are $\mathfrak{gl}_V$, $\mathfrak{sl}_V$, $\mathfrak{sp}_V$ and $\mathfrak{csp}_V$. These four cases lead to GrL , where $L = W_n, S_n, H_n$ and K_n , respectively.

In the super case the situation is much more complicated. First we consider the case of “inconsistent gradation”, meaning that $\mathfrak{g}_{-1}$ contains a non-zero even element. The classification of such strongly transitive modules is rather long and the answer consists of a dozen series and a half dozen exceptions (see [K7], Theorem 3.1). Using similar restrictions on $\mathfrak{g}_{-2}, \mathfrak{g}_{-3}, \dots$, we obtain a complete list of possibilities for $GrL_{\leq} := \bigoplus_{j \leq 0} \mathfrak{g}_j$, in the case when $\mathfrak{g}_{-1}$ contains non-zero even elements. It turns out that all but one exception are not exceptions at all, but correspond to the beginning members of some series. As a result, only $E(4|4)$ “survives”.

Step 4. Next, we turn to the case of a consistent gradation, i.e., when $\mathfrak{g}_{-1}$ is purely odd. But then $\mathfrak{g}_0$ is an “honest” Lie algebra, having a faithful irreducible representation in $\mathfrak{g}_{-1}$ (condition (G4) becomes vacuous). An explicit description of such representations is given by the classical Cartan-Jacobson theorem. In this case I use the “growth” method developed in [K1] and [K2] to determine a complete list of possibilities for $GrL_{\leq}$. This case produces mainly the (remaining four) exceptions.

Step 5 is rather long and tedious [CK3]. For each $GrL_{\leq}$ obtained in Steps 3 and 4 we determine all possible “prolongations”, i.e., infinite-dimensional $\mathbb{Z}$ -graded Lie superalgebras satisfying (G2), whose negative part is the given $GrL_{\leq}$.

Step 6. It remains to reconstruct L from GrL , i.e., to find all possible filtered simple linearly compact Lie superalgebras L with given GrL (such an L is called a simple filtered deformation of GrL). Of course, there is a trivial filtered deformation: $GrL := \Pi_{j \geq -d} \mathfrak{g}_j$, which is simple iff GrL is. It is proved in [CK2] by a long and tedious calculation that only $SHO(m|m)$ for m even ≥ 2 and $SKO(m|m+1; \frac{m+2}{m})$ for m odd ≥ 3 have a non-trivial simple filtered deformation, which are the ninth and tenth series. It would be nice to have a more conceptual proof. Recall that $SHO(m|m)$ is not simple, though it does have a simple filtered deformation. Note also that in the Lie algebra case all filtered deformations are trivial.

1.5. Plan of the proof of Theorem 2.

The key idea is the same as in the proof of Theorem 1. Choose a maximal subalgebra L_0 of a simple finite-dimensional Lie superalgebra L containing the even part of L . It is easy to see then that $L_{-1} = L$, so that the corresponding Weisfeiler’s filtration has depth 1. Hence GrL has the form: $GrL = \bigoplus_{j=-1}^N \mathfrak{g}_j$. This gradation is consistent and, of course, satisfies conditions (G0)–(G3). There are two cases.

Case 1. $N \geq 1$. Then we apply the growth method (as in Step 4 of Sec. 1.4.) to obtain a complete list of possibilities for $GrL_{\leq}$. Then, as in Step 5 of Sec. 1.4. we determine all prolongations of each $GrL_{\leq}$ (all of them will be subalgebras of $W(0, \dim \mathfrak{g}_1)$), and all filtered deformations of these prolongations. This case produces all “non-classical” series, and also $sl(m|n)$, $spo(m|2)$ and $p(m|m)$.

Case 2. $N = 0$. Then $L_{\overline{0}}$ is a semisimple Lie algebra and its representation in $L_{\overline{1}}$ is irreducible. The Killing form on L is either non-degenerate, in which case we apply the standard Killing-Cartan techniques, or it is identically zero. In the latter case one uses Dynkin’s index to find all possibilities for the $L_{\overline{0}}$ -module $L_{\overline{1}}$.

1.6. In order to describe the construction of the exceptional infinite-dimensional Lie superalgebras (given in [CK3]), I need to make some remarks. Let $\Omega_m = \Omega(m|0)$ be the algebra of differential forms over $\mathcal{O}_m$, let Ω_m^k denote the space of forms of degree k , and $\Omega_{m,cl}^k$ the subspace of closed forms. For any $\lambda \in \mathbb{C}$ the representation of W_m on Ω_m^k can be “twisted” by letting

$$X \mapsto L_X + \lambda \operatorname{div} X, \quad X \in W_m,$$

to get a new W_m -module, denoted by $\Omega_m^k(\lambda)$ (the same can be done for $W(m|n)$). Obviously, $\Omega_m^k(\lambda) = \Omega_m^k$ when restricted to S_m . Then we have the following obvious W_m -module isomorphisms: $\Omega_m^0 \simeq \Omega_m^m(-1)$ and $\Omega_m^0(1) \simeq \Omega_m^m$. Furthermore, the map $X \mapsto \iota_X(dx_1 \wedge \dots \wedge dx_m)$ gives the following W_m -module and S_m -module isomorphisms:

$$W_m \simeq \Omega_m^{m-1}(-1), \quad S_m \simeq \Omega_{m,cl}^{m-1}.$$

We shall identify the representation spaces via these isomorphisms.

The simplest is the construction of the largest exceptional Lie superalgebra $E(5|10)$. Its even part is the Lie algebra S_5 , its odd part is the space of closed 2-forms $\Omega_{5,cl}^2$. The remaining commutators are defined as follows for $X \in S_5$, $\omega, \omega' \in \Omega_{5,cl}^2$:

$$[X, \omega] = L_X \omega, \quad [\omega, \omega'] = \omega \wedge \omega' \in \Omega_{5,cl}^4 = S_5.$$

Each quintuple of integers $(a_1, a_2, \dots, a_5)$ such that $a = \sum_i a_i$ is even, defines a $\mathbb{Z}$ -gradation of $E(5|10)$ by letting:

$$\deg x_i = -\frac{\partial}{\partial x_i} = a_i, \quad \deg dx_i = a_i - \frac{1}{4}a.$$

The quintuple $(2, 2, \dots, 2)$ defines the (only) consistent $\mathbb{Z}$ -gradation, which has depth 2: $E(5|10) = \Pi_{j \geq -2} \mathfrak{g}_j$, and one has:

$$\mathfrak{g}_0 \simeq sl_5 \text{ and } \mathfrak{g}_{-1} \simeq \Lambda^2 \mathbb{C}^5, \quad \mathfrak{g}_{-2} \simeq \mathbb{C}^{5*} \text{ as } \mathfrak{g}_0\text{-modules.}$$

Furthermore, $\Pi_{j \geq 0} \mathfrak{g}_j$ is a maximal open subalgebra of $E(5|10)$ (the only one which is invariant with respect to all automorphisms). There are three other maximal open subalgebras in $E(5|10)$, associated to $\mathbb{Z}$ -gradations corresponding to quintuples $(1, 1, 1, 1, 2)$, $(2, 2, 2, 1, 1)$ and $(3, 3, 2, 2, 2)$, and one can show that these four are all, up to conjugacy, maximal open subalgebras (cf. [CK3]).

Another important $\mathbb{Z}$ -gradation of $E(5|10)$, which is, unlike the previous four, by infinite-dimensional subspaces, corresponds to the quintuple $(0, 0, 0, 1, 1)$ and has depth 1: $E(5|10) = \Pi_{\lambda \geq -1} \mathfrak{g}^\lambda$. One has: $\mathfrak{g}^0 \simeq E(3|6)$ and the $\mathfrak{g}^\lambda$ form an important family of irreducible $E(3|6)$ -modules [KR2]. The consistent $\mathbb{Z}$ -gradation of $E(5|10)$ induces that of $\mathfrak{g}^0 : E(3|6) = \Pi_{j \geq -2} \mathfrak{a}_j$, where

$$\mathfrak{a}_0 \simeq sl_3 \oplus sl_2 \oplus gl_1, \quad \mathfrak{a}_{-1} \simeq \mathbb{C}^3 \boxtimes \mathbb{C}^2 \boxtimes \mathbb{C}, \quad \mathfrak{a}_{-2} \simeq \mathbb{C}^3 \boxtimes \mathbb{C} \boxtimes \mathbb{C}.$$

A more explicit construction of $E(3|6)$ is as follows [CK3]: the even part is $W_3 + \Omega_3^0 \otimes sl_2$, the odd part is $\Omega_3^1(-\frac{1}{2}) \otimes \mathbb{C}^2$ with the obvious action of the even part, and the bracket of two odd elements is defined as follows:

$$[\omega \otimes u, \omega' \otimes v] = (\omega \wedge \omega') \otimes (u \wedge v) + (d\omega \wedge \omega' + \omega \wedge d\omega') \otimes (u \cdot v).$$

Here the identifications $\Omega_3^2(-1) = W_3$ and $\Omega_3^0 = \Omega_3^3(-1)$ are used.

The gradation of $E(5|10)$ corresponding to the quintuple $(0, 1, 1, 1, 1)$ has depth 1 and its 0th component is isomorphic to $E(1|6)$ (cf. [CK3]).

The construction of $E(4|4)$ is also very simple [CK3]: The even part is W_4 , the odd part is $\Omega_4^1(-\frac{1}{2})$ and the bracket of two odd elements is:

$$[\omega, \omega'] = d\omega \wedge \omega' + \omega \wedge d\omega' \in \Omega_4^3(-1) = W_4.$$

The construction of $E(3|8)$ is slightly more complicated, and we refer to [CK3] for details.

1.7. All exceptional simple finite-dimensional Lie superalgebras (including the exceptional Lie algebras) are obtained as special cases of the following important construction [K2]. Let $I = \{1, \dots, r\}$ and let $I_{\overline{1}}$ be a subset of I , $I_{\overline{0}} = I \setminus I_{\overline{1}}$. Let $A = (a_{ij})_{i,j \in I}$ be a matrix over $\mathbb{C}$. We associate to the pair $(A, I_{\overline{1}})$ a Lie superalgebra $\mathfrak{g}(A, I_{\overline{1}})$ as follows. Let $\tilde{\mathfrak{g}}(A, I_{\overline{1}})$ be the Lie superalgebra on generators e_i, f_i, h_i ($i \in I$) of parity $p(h_i) = \overline{0}$ for $i \in I$, $p(e_i) = p(f_i) = \alpha \in \{\overline{0}, \overline{1}\}$ for $i \in I_\alpha$, and the following standard relations:

$$[h_i, h_j] = 0, [e_i, f_j] = \delta_{ij} h_i, [h_i, e_j] = a_{ij} e_j, [h_i, f_j] = -a_{ij} f_j.$$

Define a $\mathbb{Z}$ -gradation $\tilde{\mathfrak{g}}(A, I_{\overline{1}}) = \bigoplus_{j \in \mathbb{Z}} \tilde{\mathfrak{g}}_j$ by letting $\deg h_i = 0$, $\deg e_i = -\deg f_i = 1$. Then $\tilde{\mathfrak{g}}_0$ is the $\mathbb{C}$ -span of $\{h_i\}_{i \in I}$, and we denote by $J(A, I_{\overline{1}})$ the sum of all $\mathbb{Z}$ -graded ideals of $\tilde{\mathfrak{g}}(A, I_{\overline{1}})$ that intersect $\tilde{\mathfrak{g}}_0$ trivially. We let

$$\mathfrak{g}(A, I_{\overline{1}}) = \tilde{\mathfrak{g}}(A, I_{\overline{1}}) / J(A, I_{\overline{1}}).$$

Of course, if A is the Cartan matrix of a simple finite-dimensional Lie algebra $\mathfrak{g}$, then $\mathfrak{g} \simeq \mathfrak{g}(A, \emptyset)$, the ideal $J(A, \emptyset)$ being generated by “Serre relations”. Likewise, generalized Cartan matrices give rise to Kac-Moody Lie algebras [K3].

Consider the following matrices ($a \in \mathbb{C} \setminus \{0, -1\}$):

$$D_a = \begin{bmatrix} 0 & -1 & -a \\ -1 & 2 & 0 \\ -1 & 0 & 2 \end{bmatrix}, \quad F = \begin{bmatrix} 0 & -1 & 0 & 0 \\ -1 & 2 & -2 & 0 \\ 0 & -1 & 2 & -1 \\ 0 & 0 & -1 & 2 \end{bmatrix}, \quad G = \begin{bmatrix} 0 & -1 & 0 \\ -1 & 2 & -3 \\ 0 & -1 & 2 \end{bmatrix}.$$

Then $D(2, 1; \alpha) \simeq \mathfrak{g}(D_\alpha, \{1\})$, $F(4) \simeq \mathfrak{g}(F, \{1\})$ and $G(3) \simeq \mathfrak{g}(G, \{1\})$. Note, however, that unlike in the Lie algebra case, “inequivalent” pairs (A, I_Γ) may produce isomorphic Lie superalgebras. For example, in the cases $D(2, 1; a)$, $F(4)$ and $G(3)$ there are 2, 6 and 4 such pairs, respectively.

Finite-dimensional simple Lie superalgebras that are isomorphic to $\mathfrak{g}(A, I_\Gamma)$ for some matrix A are called *basic* (they will play an important role in the next parts of the talk). The remaining basic simple Lie superalgebras (that are not Lie algebras) are $sl(m|n)/\delta_{m,n}\mathbb{C}E$ and $spo(m|n)$.

2. A classification of superconformal algebras

Superconformal algebras have been playing an important role in superstring theory and in conformal field theory. Here I will explain how to apply Theorem 1 to the classification of “linear” superconformal algebras. By a (“linear”) superconformal algebra I mean a Lie superalgebra $\mathfrak{g}$ spanned by coefficients of a finite collection F of fields such that the following two properties hold:

- (1) for $a, b \in F$ the singular part of OPE is finite, i.e.,

$$[a(z), b(w)] = \sum_j c_j(w) \partial_w^j \delta(z-w) \quad (\text{a finite sum}), \text{ where all } c_j(w) \in \mathbb{C}[\partial_w]F,$$

- (2) $\mathfrak{g}$ contains no non-trivial ideals spanned by coefficients of fields from a $\mathbb{C}[\partial_w]$ -submodule of $\mathbb{C}[\partial_w]F$.

(Recall that a field is a formal expression $a(z) = \sum_{n \in \mathbb{Z}} a_n z^n$, where $a_n \in \mathfrak{g}$ and z is an indeterminate, and $\delta(z-w) = z^{-1} \sum_{n \in \mathbb{Z}} (w/z)^n$ is the formal δ -function. See [K5] for details.)

This problem goes back to the physics paper [RS], some progress in its solution was made in [K6] and a complete solution was given in [FK]. (A complete classification even in the “quadratic” case seems to be a much harder problem, see [FL] and Section 4 below for some very interesting examples.) The simplest example is the loop algebra $\tilde{\mathfrak{g}} = \mathbb{C}[x, x^{-1}] \otimes \mathfrak{g}$ (= centerless affine Kac-Moody superalgebra), where $\mathfrak{g}$ is a simple finite-dimensional Lie superalgebra. Then $F = \{a(z) = \sum_{n \in \mathbb{Z}} (x^n \otimes a) z^{-n-1}\}_{a \in \mathfrak{g}}$, and $[a(z), b(w)] = [a, b](w) \delta(z-w)$. The next example is the Lie algebra $\text{Vect } \mathbb{C}^\times$ of regular vector fields on $\mathbb{C}^\times$ (= centerless Virasoro algebra); F consists of one field, the Virasoro field $L(z) = -\sum_{n \in \mathbb{Z}} (x^n \frac{d}{dx}) z^{-n-1}$, and $[L(z), L(w)] = \partial_w L(w) \delta(z-w) + 2L(w) \delta'_w(z-w)$.

One of the main theorems of [DK] states that these are all examples in the Lie algebra case. The strategy of the proof is the following. Let $\partial = \partial_z$ and consider the (finitely generated) $\mathbb{C}[\partial]$ -module $R = \mathbb{C}[\partial]F$. Define the “ λ -bracket” $R \otimes R \rightarrow \mathbb{C}[\lambda] \otimes R$ by the formula: $[a_\lambda b] = \sum_j \lambda^j c_j$. This satisfies the axioms of a *conformal (super)algebra* (see [DK], [K5]), similar to the Lie (super)algebra axioms:

- (i) $[\partial a_\lambda b] = -\lambda[a_\lambda b]$, $[a_\lambda \partial b] = (\partial + \lambda)[a_\lambda b]$,
- (ii) $[a_\lambda b] = -(-1)^{p(a)P(b)}[b_{-\lambda-\partial} a]$,
- (iii) $[a_\lambda [b_\mu c]] = [[a_\lambda b]_{\lambda+\mu} c] + (-1)^{p(a)p(b)}[b_\mu [a_\lambda c]]$.

The main observation of [DK] is that a conformal (super)algebra is completely determined by the Lie (super)algebra spanned by all coefficients of negative powers of z of the fields $a(z)$ from F , called the *annihilation algebra*, along with an even surjective derivation of the annihilation algebra. Furthermore, apart from the case of current algebras, the completed annihilation algebra turns out to be an infinite-dimensional simple linearly compact Lie (super)algebra of growth 1. Since in the Lie algebra case the only such example is W_1 , the proof is finished.

In the superalgebra case the situation is much more interesting since there are many infinite-dimensional simple linearly compact Lie superalgebras of growth 1. By Theorem 1, the complete list is as follows:

$$W(1|N), \quad S'(1|N), \quad K(1|N) \quad \text{and} \quad E(1|6).$$

In all cases, except the second, there is a unique, up to conjugacy, even surjective derivation, hence a unique corresponding superconformal algebra. They are denoted by $W_{(N)}$, $K_{(N)}$ if $N \neq 4$, $K'_{(4)}$ and $CK_{(6)}$, respectively. The Lie superalgebras $W_{(N)}$ and $K_{(N)}$ are constructed in the same way as $W(1|N)$ and $K(1|N)$, except that one replaces $\mathcal{O}_1\langle N \rangle$ by $\mathbb{C}[x, x^{-1}]\langle N \rangle$. The construction of the exceptional superconformal algebra $CK_{(6)}$ is more difficult, and may be found in [CK1] or [K6]. However, $S'(1|N)$ has two families of even surjective derivations. The corresponding superconformal algebras are derived algebras of

$$S_{(N), \epsilon, a} = \{X \in W_{(N)} \mid \operatorname{div}(e^{ax}(1 + \epsilon \xi_1 \dots \xi_N)X) = 0\}, \quad a \in \mathbb{C}, \epsilon = \pm 1.$$

Thus, one obtains the following theorem.

Theorem 3. [FK] *A complete list of superconformal algebras consists of loop algebras $\hat{\mathfrak{g}}$, where $\mathfrak{g}$ is a simple finite-dimensional Lie superalgebra, and of Lie superalgebras ($N \in \mathbb{Z}_+$): $W_{(N)}$, $S'_{(N+2), \epsilon, a}$ (N even and $a = 0$ if $\epsilon = 1$), $K_{(N)}$ ($N \neq 4$), $K'_{(4)}$, and $CK_{(6)}$.*

Note that the first members of the above series are well-known superalgebras: $W_{(0)} \simeq K_{(0)}$ is the Virasoro algebra, $K_{(1)}$ is the Neveu-Schwarz algebra, $K_{(2)} \simeq W_{(1)}$ is the $N = 2$ algebra, $K_{(3)}$ is the $N = 3$ algebra, $S'_{(2)} = S'_{(2), 0, 0}$ is the $N = 4$ algebra, $K'_{(4)}$ is the big $N = 4$ algebra (all centerless). These algebras, along with $W_{(2)}$ and $CK_{(6)}$ are the only superconformal algebras for which all fields are primary with positive conformal weights [K6]. It is interesting to note that all of them are contained in $CK_{(6)}$, which consists of 32 fields, the even ones are the Virasoro fields and 15 currents that form $\tilde{so}_6$, and the odd ones are 6 and 10 fields of conformal weight $3/2$ and $1/2$, respectively. Here is the table of (some) inclusions, where in square brackets the number of fields is indicated:

$$\begin{array}{ccccc} CK_{(6)}[32] & \supset & W_{(2)}[12] & \supset & W_{(1)} = K_{(2)}[4] \supset K_{(1)}[2] \supset \operatorname{Vir}[1] \\ \cup & & \cup & & \\ K_{(3)}[8] \subset K'_{(4)}[16] & & S'_{(2), \epsilon, a}[8] & & \end{array}.$$

All of these Lie superalgebras have a unique non-trivial central extension, except for K'_4 that has three [KL] and $CK_{(6)}$ that has none. All other Lie superalgebras

listed by Theorem 2 have no non-trivial central extensions. (The presence of a central term is necessary for the existence of interesting representations and the construction of an interesting conformal field theory.)

3. Representations of affine superalgebras and “almost” modular forms

3.1. Finite-dimensional irreducible representations of simple finite-dimensional Lie superalgebras are much less understood than in the Lie algebra case, the main reason being the occurrence of isotropic roots in the super case. (A review may be found in the proceedings of the last ICM, see [Se].) The natural analogues of these representations in the case of affine (super)algebras are the integrable highest weight modules.

Let us first recall the basic definitions in the Lie algebra case, i.e., for an affine Kac-Moody algebra $\hat{\mathfrak{g}}$ [K3]. Let $\mathfrak{g}$ be a finite-dimensional simple Lie algebra and let $(\cdot|\cdot)$ be an invariant symmetric bilinear form on $\mathfrak{g}$ normalized by the condition that $(\alpha|\alpha) = 2$ for a long root α ($(a|b) = \text{tr}ab$ in the case $\mathfrak{g} = \mathfrak{sl}_m$). Recall that the associated *affine algebra* is

$$\hat{\mathfrak{g}} = (\mathbb{C}[x, x^{-1}] \otimes_{\mathbb{C}} \mathfrak{g}) \oplus \mathbb{C}K \oplus \mathbb{C}D$$

with the following commutation relations ($a, b \in \mathfrak{g}$; $m, n \in \mathbb{Z}$ and $a(m)$ stands for $x^m \otimes a$):

$$[a(m), b(n)] = [a, b](m+n) + m\delta_{m,-n}(a|b)K, [D, a(m)] = ma(m), [K, \hat{\mathfrak{g}}] = 0.$$

Note that the derived algebra $\hat{\mathfrak{g}}'$ is a central extension (by $\mathbb{C}K$) of the loop algebra $\tilde{\mathfrak{g}}$ that has made an appearance in Section 2. (It is also isomorphic to $\mathfrak{g}(\hat{A})$, where $\hat{A}$ is the extended Cartan matrix of $\mathfrak{g}$, cf. Sec. 1.7..) As we shall see, without central extension one loses all interesting representations. In any irreducible $\hat{\mathfrak{g}}$ -module V one has: $K = kI_V$; the number k is called the *level* of V . The scaling element D is necessary for the convergence of characters.

Choose a Cartan subalgebra $\mathfrak{h}$ of $\mathfrak{g}$ and let $\mathfrak{g} = \mathfrak{h} \oplus (\oplus_{\alpha \in \Delta} \mathfrak{g}_{\alpha})$ be the root space decomposition, where $\mathfrak{g}_{\alpha}$ denotes the root space attached to a root $\alpha \in \Delta \subset \mathfrak{h}^*$. Let $\hat{\mathfrak{h}} = \mathfrak{h} + \mathbb{C}K + \mathbb{C}D$ be the Cartan subalgebra of $\hat{\mathfrak{g}}$, and, as before, let $\mathfrak{g}_{\alpha}(m) = x^m \otimes \mathfrak{g}_{\alpha}$. We extend the invariant bilinear form from $\mathfrak{h}$ to a symmetric bilinear form on $\hat{\mathfrak{h}}$ by letting $(\mathfrak{h}|\mathbb{C}K + \mathbb{C}D) = 0$, $(K|K) = (D|D) = 0$, $(K|D) = 1$, and identify $\hat{\mathfrak{h}}$ with $\hat{\mathfrak{h}}^*$ via this form.

A $\hat{\mathfrak{g}}$ -module V is called *integrable* if the following two properties hold:

- (M1) $\hat{\mathfrak{h}}$ is diagonalizable on V ,
- (M2) for each $\alpha \in \Delta$ and $m \in \mathbb{Z}$, $\mathfrak{g}_{\alpha}(m)$ is locally finite on V .

Choose a set of positive roots $\Delta^+ \subset \Delta$, and let $\mathfrak{n}^+ = \oplus_{\alpha \in \Delta^+} \mathfrak{g}_{\alpha}$, $\hat{\mathfrak{n}}^+ = \mathfrak{n}^+ + \sum_{n \geq 1} x^n \otimes \mathfrak{g}$. For each $\Lambda \in \hat{\mathfrak{h}}^*$ one defines an *irreducible highest weight module*

$L(\Lambda)$ over $\hat{\mathfrak{g}}$ as the (unique) irreducible $\hat{\mathfrak{g}}$ -module for which there exists a non-zero vector v_Λ such that

$$hv_\Lambda = \Lambda(h)v_\Lambda \text{ for all } h \in \hat{\mathfrak{h}}, \hat{\mathfrak{n}}_+ v_\Lambda = 0.$$

Without loss of generality we shall let $\Lambda(D) = 0$; then the spectrum of $-D$ on $L(\Lambda)$ is $\mathbb{Z}_+$.

Integrable highest weight modules over affine Lie algebras (they are automatically irreducible) attracted a lot of attention in the past few decades both of mathematicians and of physicists (some aspects of the theory are discussed in [K3], [Wa2].) Here I will only mention some relevant to the talk facts. First, the level k of such a module is a non-negative integer (and $k = 0$ iff $\dim L(\Lambda) = 1$), and there is a finite number of them for each k . One of the most remarkable properties of these modules is modular invariance, which I explain below.

3.2. Let us coordinatize $\hat{\mathfrak{h}}$ by letting

$$(\tau, z, t) = 2\pi i(z - \tau D + tK),$$

where $z \in \mathfrak{h}$, $\tau, t \in \mathbb{C}$, and define the *character* of the $\hat{\mathfrak{g}}$ -module $L(\Lambda)$ by:

$$ch_\Lambda(\tau, z, t) = \text{tr}_{L(\Lambda)} e^{2\pi i(z - \tau D + tK)}.$$

If $L(\Lambda)$ is integrable, then $ch_\Lambda(\tau, z, t)$ is a holomorphic function for $(\tau, z, t) \in \mathcal{H} \times \mathfrak{h} \times \mathbb{C}$, where $\mathcal{H} = \{\tau \in \mathbb{C} \mid \text{Im } \tau > 0\}$.

Recall the following well-known action of $SL_2(\mathbb{Z})$ on $\mathcal{H} \times \mathfrak{h} \times \mathbb{C}$:

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \cdot (\tau, z, t) = \left(\frac{a\tau + b}{c\tau + d}, \frac{z}{c\tau + d}, t - \frac{c(z|z)}{2(c\tau + d)} \right).$$

Then it turns out that there exists an explicit rational number m_Λ , called *modular anomaly* (see [K3], (12.7.5)) such that the normalized character $\chi_\Lambda = e^{2\pi i m_\Lambda \tau} ch_\Lambda$ of an integrable $L(\Lambda)$ is invariant with respect to a congruence subgroup of $SL_2(\mathbb{Z})$ (see [K3], Chapter 13). If ch_Λ has this property, one says that $L(\Lambda)$ is *modular invariant*.

3.3. It turns out that $L(\Lambda)$ is modular invariant for a much wider (than integrable) collection of Λ 's, called *admissible*, defined by the condition [KW1], [KW2], [K4]:

$$2(\Lambda + \hat{\rho}|\alpha)/(\alpha|\alpha) \in \mathbb{Q} \setminus \{0, -1, -2, \dots\} \text{ for all } \alpha \in \hat{\Delta}^+ \text{ such that } (\alpha|\alpha) \neq 0.$$

(The conjecture of Wakimoto and myself is that these are all modular invariant $L(\Lambda)$.) Here $\hat{\Delta}^+$ is the set of positive roots of $\hat{\mathfrak{g}}$ corresponding to $\hat{\mathfrak{n}}^+ : \hat{\Delta}^+ = \Delta^+ \cup \{\alpha + nK \mid \alpha \in \Delta, n \geq 1\} \cup \{nK \mid n \geq 1\}$, and $\hat{\rho} \in \hat{\mathfrak{h}}^*$ is a vector satisfying $(\hat{\rho}|\alpha_i) = \frac{1}{2}(\alpha_i|\alpha_i)$ for $i = 0, 1, \dots, r$, where $\Pi = \{\alpha_1, \dots, \alpha_r\}$ are simple roots of Δ^+ , $\hat{\Pi} = \{\alpha_0 := K - \theta\} \cup \Pi$ are simple roots of $\hat{\Delta}^+$, θ is the highest root of Δ^+ .

I shall describe explicitly the most important class of them, called *principal admissible*. Fix a positive integer u and let $\hat{\Pi}_u = \{uK - \theta\} \cup \Pi$. Let $k = v/u$ be

a rational number, such that $v \in \mathbb{Z}$ is relatively prime to u and $u(k + h^\vee) \geq h^\vee$. Here $h^\vee$ is the dual Coxeter number (defined in a more general Lie superalgebra context further on). Let W be the Weyl group of $\mathfrak{g}$ and let $P^\vee = \{\lambda \in \mathfrak{h} | (\lambda|\alpha) \in \mathbb{Z} \text{ for all } \alpha \in \Delta\}$. For each $\alpha \in P^\vee$ define a translation $t_\alpha \in \text{End } \hat{\mathfrak{h}}$ by the formula $t_\alpha(\lambda) = \lambda + (\lambda|K)\alpha - ((\lambda|\alpha) + \frac{(\lambda|K)}{2}(\alpha|\alpha))K$. Pick an element $\hat{w} = t_\beta w$, where $w \in W$, such that $\hat{w}\hat{\Pi}_u \subset \hat{\Delta}^+$ (these are all subsets of $\hat{\Delta}^+$ isomorphic to $\hat{\Pi}$). Let Λ^0 be an integrable highest weight of level $u(k + h^\vee) - h^\vee$. Then

$$\Lambda = \hat{w}(\Lambda^0 + \hat{\rho} - (u-1)(k + h^\vee)D) - \hat{\rho}$$

is a *principal admissible weight* of level k . The character of the corresponding $L(\Lambda)$ is given by the following formula [KW1]–[KW2]:

$$\left(\hat{R}ch_{L(\Lambda)} \right) (\tau, z, t) = \left(\hat{R}ch_{L(\Lambda^0)} \right) (u\tau, w^{-1}(z + \tau\beta), u^{-1}(t + (z|\beta) + \frac{1}{2}\tau(\beta|\beta))) , \quad (\text{Ch})$$

where $\hat{R} = e^{\hat{\rho}} \prod_{\alpha \in \hat{\Delta}^+} (1 - e^{-\alpha})^{\text{mult } \alpha}$ is the Weyl denominator function for $\hat{\mathfrak{g}}$, and $\text{mult } \alpha = 1$ except for $\text{mult } nK = r$ for all n (note that this formula is a tautology if $\Lambda = \Lambda^0$ is integrable; this happens iff $u = 1$). Recall that $ch_{L(\Lambda^0)}$ is given by the Weyl-Kac character formula [K3].

3.4. Let now $\mathfrak{g} = \mathfrak{g}(A)$ be a basic simple finite-dimensional Lie superalgebra (see Sec. 1.7). Then $\mathfrak{g}$ carries a unique, up to a constant factor, non-degenerate invariant bilinear form B (“invariant” means that $B([a, b], c) = B(a, [b, c])$). Let $\mathfrak{h} = \sum_{i=1}^r \mathbb{C}h_i$ be the Cartan subalgebra of $\mathfrak{g}$, $\mathfrak{n}^+$ the subalgebra of $\mathfrak{g}$ generated by all e_i , $\Delta^+ \subset \mathfrak{h}^*$ the set of positive roots (i.e., roots of $\mathfrak{h}$ in $\mathfrak{n}^+$), $\Delta = \Delta^+ \cup -\Delta^+$ the set of all roots, $\Delta_{\overline{0}}$ and $\Delta_{\overline{1}}$ the sets of even and odd roots, $\{\alpha_1, \dots, \alpha_r\} \subset \Delta^+$ the set of simple roots ($\alpha_i(h_j) = a_{ji}$), $\theta \in \Delta^+$ the highest root. Define $\rho \in \mathfrak{h}^*$ by $B(\rho, \alpha_i) = \frac{1}{2}B(\alpha_i, \alpha_i)$, $i = 1, \dots, r$. Then $h_B^\vee = B(\rho, \rho) + \frac{1}{2}B(\theta, \theta)$ is the eigenvalue of the Casimir operator in the adjoint representation.

If $h_B^\vee \neq 0$, we let $\Delta_0^\# = \{\alpha \in \Delta_{\overline{0}} | h_B^\vee B(\alpha, \alpha) > 0\}$. If $h_B^\vee = 0$, which happens for $\mathfrak{g} = \mathfrak{sl}(m|m)/\mathbb{C}E$, $\mathfrak{spo}(2m|2m+2)$ and $D(2, 1; a)$, we take for $\Delta_0^\#$ the sets of roots of the subalgebra $\mathfrak{sl}_m$, $\mathfrak{so}_{2m+2}$ and $\mathfrak{sl}_2 \oplus \mathfrak{sl}_2$, respectively. Let $W^\#$ denote the subgroup of the Weyl group of $\mathfrak{g}_{\overline{0}}$ generated by reflections with respect to all $\alpha \in \Delta_0^\#$. Denote by $(\cdot | \cdot)$ the invariant bilinear form on $\mathfrak{g}$ normalized by the condition $(\alpha|\alpha) = 2$ for the longest root $\alpha \in \Delta_0^\#$. The corresponding to this form number $h^\vee = h_{(\cdot | \cdot)}^\vee$ is called the *dual Coxeter number*. (For example, this number equals $|m - n|$ for $\mathfrak{sl}(m|n)$, $\frac{1}{2}(m - n) + 1$ for $\mathfrak{spo}(m|n)$ with $m \geq n - 2$, 30 for E_8 , 3 for $F(4)$, 2 for $G(3)$.)

3.5. Define the affine superalgebra $\hat{\mathfrak{g}}$ associated to the Lie superalgebra $\mathfrak{g}$ in exactly the same way as in the Lie algebra case. The highest weight $\hat{\mathfrak{g}}$ -modules $L(\Lambda)$ are defined in the same way too. The *integrability* of a $\hat{\mathfrak{g}}$ -module V is defined by (M1), (M2) with Δ replaced by $\Delta_0^\#$, and

(M3) V is locally $\mathfrak{g}$ -finite. ((M1) and (M2) imply (M3), but there are no integrable $L(\Lambda)$ in the super case if one doesn't replace Δ by $\Delta_0^\#$.)

Integrable $\hat{\mathfrak{g}}$ -modules $L(\Lambda)$ were classified in [KW4]. However, very little is known about their characters. The following example shows that modular invariance fails already in the simplest case $\mathfrak{g} = \mathfrak{sl}(2|1)$, $\Lambda = D$. In this case

$$e^{-D} ch_{L(D)} = \prod_{n=1}^{\infty} ((1 - q^n)^{-1})(1 + z_1 q^n)(1 + z_1^{-1} q^{n-1}) A(z_1^{-1}, z_2 q^{1/2}, q), \quad (\text{A})$$

where $z_i = e^{\epsilon_i + \epsilon_3}$ ($i = 1, 2$; ϵ_i is the standard basis of the space of 3×3 diagonal matrices), $q = e^{2\pi i \tau}$, and

$$A(x, z, q) = \sum_{n \in \mathbb{Z}} \frac{q^{n^2/2} z^n}{1 + x q^n}.$$

The function $A(x, z, q)$ converges to a meromorphic function in the domain $x, y, z \in \mathbb{C}$, $|q| < 1$, and is called Appell's function. Since the first factor in (A) has the modular invariance property and $A(x, z, q)$ doesn't have it, we see that $L(D)$ is not modular invariant. We call the Appell function an *almost modular form* since it is a section of a rank 2 vector bundle on an elliptic curve for each $\tau \in \mathcal{H}$ [P] (whereas modular forms are sections of rank 1 vector bundles on it).

We call a weight $\Lambda \in \hat{\mathfrak{h}}^*$ of the Lie superalgebra $\hat{\mathfrak{g}}$ *admissible* (resp. *principal admissible*) if Λ is admissible (resp. principle admissible) for the affine Lie algebra associated to a semisimple Lie algebra with root system $\Delta_0^\#$, and we conjecture [KW4] that formula (Ch) still holds, where $\hat{R}$ in the super case is defined by $\hat{R} = e^{\hat{\rho}} \prod_{\alpha \in \hat{\Delta}_+} (1 - (-1)^{p(\alpha)} e^{-\alpha})^{(-1)^{p(\alpha)} \text{mult } \alpha}$. Unfortunately, $ch_{L(\Lambda^0)}$ is not known in general, but in the *boundary level* case, i.e., when $u(k + h^\vee) = h^\vee$, the level of Λ^0 is zero, hence $ch_{L(\Lambda^0)} = 1$, and formula (Ch) gives an explicit expression for $ch_{L(\Lambda)}$. In particular, the character is modular invariant in this case.

Let me mention in conclusion of this section that the Weyl-Kac character formula in the case of 1-dimensional module over an affine Lie algebra $\hat{\mathfrak{g}}$ turns into celebrated Macdonald's identities that express $\hat{R}$ as an infinite series, the special case for $\mathfrak{g} = \mathfrak{sl}_2$ being the Jacobi triple product identity. A sum formula for $\hat{R}$ in the super case is also known [KW3] (see also the talk [Wa1] at the last ICM). In the simplest case of $\mathfrak{g} = \mathfrak{sl}(2|1)$ one gets the identity:

$$\prod_{n=1}^{\infty} \frac{(1 - q^n)^2 (1 - uvq^{n-1})(1 - u^{-1}v^{-1}q^n)}{(1 - uq^{n-1})(1 - u^{-1}q^n)(1 - vq^{n-1})(1 - v^{-1}q^n)} = \left(\sum_{m,n=0}^{\infty} - \sum_{m,n=-1}^{-\infty} \right) u^m v^n q^{mn},$$

which goes back to Ramanujan and even further back to Kronecker.

4. Quantum reduction for affine Lie superalgebras

4.1. This part of my talk is based on a joint work with S.-S. Roan and M. Wakimoto [KRW], [KW5]. I will explain a general quantum reduction scheme, which is a further development of a number of works. They include [DS], [KS] and [Kh] on classical Drinfeld-Sokolov reduction, and [FF1], [FF2], [FKW], [B], [BT] on its quantization. As in [FF2], the basic idea is to translate the geometric Drinfeld-Sokolov reduction to a homological language as in [KS], and then to quantize this homology complex. Remarkably, this procedure gives, starting from affine superalgebras, a number of very interesting super extensions of the Virasoro algebra and their most interesting representations. I will use the very convenient language of vertex algebras (introduced in [B]), all related notations can be found in [K5].

4.2. Let $\mathfrak{g}$ be a basic simple finite-dimensional Lie superalgebra with a non-degenerate invariant bilinear form $(\cdot | \cdot)$. Fix a number k and a nilpotent even element f of $\mathfrak{g}$. Include f in an sl_2 -triple $\{e, h, f\}$ so that $[h, e] = 2e$, $[h, f] = -2f$, $[e, f] = h$. We have the eigenspace decomposition of $\mathfrak{g}$ with respect to $\text{ad } h$: $\mathfrak{g} = \bigoplus_{j \in \mathbb{Z}} \mathfrak{g}_j$, and we let $\mathfrak{g}_+ = \bigoplus_{j > 0} \mathfrak{g}_j$. The element f defines a non-degenerate skew-supersymmetric bilinear form $\langle \cdot, \cdot \rangle$ on $\mathfrak{g}_1$ by the formula $\langle a, b \rangle = (f | [a, b])$. Let A_{ne} denote the superspace $\mathfrak{g}_1$ with the form $\langle \cdot, \cdot \rangle$. Denote by A_{ch} the superspace $\pi \mathfrak{g}_+ + \pi \mathfrak{g}_+^*$, where π stands for the reversal of parity, with the skew-supersymmetric bilinear form defined by $(a, b^*) = b^*(a)$ for $a \in \pi \mathfrak{g}_+$, $b \in \pi \mathfrak{g}_+^*$, $(\pi \mathfrak{g}_+, \pi \mathfrak{g}_+) = 0 = (\pi \mathfrak{g}_+^*, \pi \mathfrak{g}_+^*)$.

Let $V^k(\hat{\mathfrak{g}})$ be the universal affine vertex algebra, and let $F^1(A_{ne})$, $F^1(A_{ch})$ be the free fermionic vertex algebras ([K5], § 4.7). Consider the vertex algebra

$$C(\mathfrak{g}, f, k) = V^k(\hat{\mathfrak{g}}) \otimes F^1(A_{ne}) \otimes F^1(A_{ch}),$$

and define its charge decomposition $C(\mathfrak{g}, f, k) = \bigoplus_{m \in \mathbb{Z}} C_m$ by letting charge $V^k(\hat{\mathfrak{g}}) =$ charge $F^1(A_{ne}) = 0$, charge $\pi \mathfrak{g}_+ = 1 = -$ charge $\pi \mathfrak{g}_+^*$.

Next, we define a differential d on $C(\mathfrak{g}, f, k)$ which makes it a homology complex. For this choose a basis $\{u_i\}_{i \in S'}$ of $\mathfrak{g}_1$ and extend it to a basis $\{u_i\}_{i \in S}$ of $\mathfrak{g}_+$ compatible with its $\mathbb{Z}$ -gradation. Denote by $\{\varphi_i\}_{i \in S}$ and $\{\varphi_i^*\}_{i \in S}$ the corresponding dual bases of $\pi \mathfrak{g}_+$ and $\pi \mathfrak{g}_+^*$, and by $\{\Phi_i\}_{i \in S'}$ the corresponding basis of A_{ne} . Consider the following odd field of the vertex algebra $C(\mathfrak{g}, f, k)$:

$$\begin{aligned} d(z) &= \sum_{i \in S} (-1)^{p(u_i)} u_i(z) \otimes \varphi_i^*(z) \otimes 1 \\ &\quad - \frac{1}{2} \sum_{i, j, k \in S} (-1)^{p(u_i)p(u_k)} c_{ij}^k \otimes \varphi_k(z) \varphi_i^*(z) \varphi_j^*(z) \otimes 1 \\ &\quad + \sum_{i \in S} (f | u_i) \otimes \varphi_i^*(z) \otimes 1 + \sum_{i \in S'} 1 \otimes \varphi_i^*(z) \otimes \Phi_i(z), \end{aligned}$$

where $[u_i, u_j] = \sum_k c_{ij}^k u_k$ in $\mathfrak{g}_+$, and let $d = \text{Res}_{z=0} d(z)$. (Note that the first two summands of d form the usual differential of a Lie (super)algebra complex.) Then one checks that $[d(z), d(w)] = 0$, hence $d^2 = 0$. It is also clear that $dC_m \subset C_{m-1}$. We define the vertex algebra $W(\mathfrak{g}, f, k)$ as the 0th homology of this complex, and call it the *quantum reduction* of the triple $(\mathfrak{g}, f, k)$ (actually, it depends only on $\mathfrak{g}, k$ and the conjugacy class of f in $\mathfrak{g}_0$).

One of the fields of the vertex algebra $W(\mathfrak{g}, f, k)$ is the following Virasoro field

$$L(z) = \frac{1}{2(k+h^\vee)} \sum_i : a_i(z) b_i(z) : + \frac{1}{2} \partial_z h(z) + \sum_{i \in S} ((1-m_i) : \partial_z \varphi_i^*(z) \varphi_i(z) : - m_i : \varphi_i^*(z) \partial_z \varphi_i(z) :) + \frac{1}{2} \sum_{i \in S'} g^{ij} : \partial_z \Phi_i(z) \Phi_j(z) :,$$

where $\{b_i\}$ and $\{a_i\}$ are dual bases of $\mathfrak{g}$: $(b_i|a_j) = \delta_{ij}$, $[h, u_i] = 2m_i u_i$, (g^{ij}) is the matrix inverse to $(\langle u_i, u_j \rangle)_{i,j \in S'}$. The central charge of $L(z)$ is equal to:

$$c(k) = \frac{k \operatorname{sdim} \mathfrak{g}}{k+h^\vee} - 3(h|h)k - \sum_{i \in S} (-1)^{p(u_i)} (12m_i^2 - 12m_i + 2) - \frac{1}{2} \operatorname{sdim} \mathfrak{g}_1.$$

Here $\operatorname{sdim} V = \dim V_0 - \dim V_1$ is the superdimension of the superspace V . Thus, all $W(\mathfrak{g}, f, k)$ are super-extensions of the Virasoro algebra. Furthermore, for each ad h eigenvector with eigenvalue $-2j$ in the centralizer of f in $\mathfrak{g}$, $W(\mathfrak{g}, f, k)$ contains a field of conformal weight $1+j$ (so that $L(z)$ corresponds to f), and these fields generate the vertex algebra $W(\mathfrak{g}, f, k)$.

Examples. (1) $\mathfrak{g}$ is a simple Lie algebra.

(a) f is a principal nilpotent element. Then $W(\mathfrak{g}, f, k)$ is called the quantum Drinfeld-Sokolov reduction. These algebras and their representations were extensively studied in [FF2], [FB], [FKW] and many other papers. The simplest case of $\mathfrak{g} = \mathfrak{sl}_2$ produces the Virasoro vertex algebra. The case $\mathfrak{g} = \mathfrak{sl}_3$ gives the W_3 algebra [Z].

(b) f is a lowest root vector of $\mathfrak{g}$. These vertex algebras were discussed from a different point of view in [FL] under the name quasi-superconformal algebras. The special case of $\mathfrak{g} = \mathfrak{sl}_3$ was studied from a quantum reduction viewpoint in [Be].

(2) $\mathfrak{g}$ is a simple basic Lie superalgebra and f is an even lowest root vector.

(a) One has the following correspondence:

$\mathfrak{g}$	$\mathfrak{spo}(2 1)$	$\mathfrak{sl}(2 1)$	$\mathfrak{sl}(2 2)$	$\mathfrak{spo}(2 3)$	$D(2, 1; a)$
$W(\mathfrak{g}, f, k)$	Neveu-Schwarz	$N = 2$	$N = 4$	$N = 3$	big $N = 4$

(In the last two columns one gets an isomorphism after adding one fermion, resp. four fermions and one boson.).

(b) Almost every lowest root vector of a simple component of $\mathfrak{g}_0$ can be made equal f . This gives all superconformal algebras of [FL] (by definition, they are generated by the Virasoro field, the even fields of weight 1 and N odd fields of weight $3/2$), and many new examples.

4.3. Let M be a highest weight module over $\hat{\mathfrak{g}}$. It extends to a vertex algebra module over $V^k(\hat{\mathfrak{g}})$, and we consider the $C(\mathfrak{g}, f, k)$ -module $C(M) = M \otimes F^1(A_{ne}) \otimes F^1(A_{ch})$. The element d acts on $C(M)$ and again $d^2 = 0$, hence we can consider homology $H(M) = \bigoplus_j H_j(M)$ which is a module over $W(\mathfrak{g}, f, k)$. The $W(\mathfrak{g}, f, k)$ -module $H(M)$ is called the *quantum reduction* of the $\hat{\mathfrak{g}}$ -module M . Using the Euler-Poincaré principle one easily computes the character of $H(M)$ in terms of ch_M . The

basic conjecture of [FKW], [KRW] is that $H(M)$ is irreducible (in particular, at most one $H_j(M)$ is non-zero) if M is admissible.

Examples. (1) $\mathfrak{g} = \mathfrak{sl}_2$. Let k be an admissible level, i.e., k is a rational number with positive denominator u such that $u(k+2) \geq 2$ (recall that $h^\vee = 2$). The set of principal admissible weights of level k is as follows (α is a simple root of $\mathfrak{sl}_2$) [KW1], [K4]:

$$\{\Lambda_{k,j,n} = kD + \frac{1}{2}(n - j(k+2))\alpha \mid 0 \leq j \leq u-1, 0 \leq n \leq u(k+2) - 2\}.$$

Then the quantum reduction of the $\hat{\mathfrak{sl}}_2$ -module $L(\Lambda_{k,j,n})$ is the “minimal series” module corresponding to parameters $p = u(k+2)$, $p' = u$ (cf. [BPZ], [K4]):

$$c^{(p,p')} = 1 - 6 \frac{(p-p')^2}{pp'}, \quad h_{j+1,n+1}^{(p,p')} = \frac{(p(j+1) - p'(n+1))^2 - (p-p')^2}{4pp'}.$$

The character formula (Ch) for $L(\Lambda_{k,j,n})$ gives immediately all the characters of minimal series.

(2) $\mathfrak{g} = \mathfrak{spo}(2|1)$. We get all minimal series modules over the Neveu-Schwarz algebra and their characters by quantum reduction of all (not only principal) admissible $\hat{\mathfrak{g}}$ -modules.

(3) $\mathfrak{g} = \mathfrak{sl}(2|1)$. Then the boundary admissible levels are $k = m^{-1} - 1$, where $m \in \mathbb{Z}$, $m \geq 2$. One has the following $\hat{w}\hat{\Pi}_m$'s (see Sec. 3.3.), where α_1 and α_2 are odd: $\{\alpha_0 + b_0K, \alpha_1 + b_1K, \alpha_2 + b_2K\}$. Here b_i are non-negative integers, $b_0 \geq 1$ and $\sum b_i = m - 1$. The quantum reduction of the corresponding admissible $\hat{\mathfrak{g}}$ -modules gives all the minimal series representations of the $N = 2$ superconformal algebra (cf. [FST] and references there). Again, formula (Ch) gives immediately their characters.

(4) $\mathfrak{g} = \mathfrak{sl}(2|2)$ (resp. $\mathfrak{spo}(2|3)$). In a similar fashion we recover the characters of $N = 4$ [ET] (resp. $N = 3$ [M]) superconformal algebras.

5. Representations of $E(3|6)$ and the standard model

5.1. By a representation of a linearly compact Lie superalgebra L we shall mean a continuous representation in a vector space V with discrete topology (then the contragredient representation is a continuous representation in a linearly compact space V^*). Fix an open subalgebra L_0 of L . We shall assume that V is locally L_0 -finite, meaning that any vector of V is contained in a finite-dimensional L_0 -invariant subspace (this property actually often implies that V is continuous). These kinds of representations were studied in the Lie algebra case by Rudakov [R]. It is easy to show that such an irreducible L -module V is a quotient of an induced module $\text{Ind}_{L_0}^L U = U(L) \otimes_{U(L_0)} U$, where U is a finite-dimensional irreducible L_0 -module, by a (unique in good cases) maximal submodule. The induced module $\text{Ind}_{L_0}^L U$ is called *degenerate* if it is not irreducible. An irreducible quotient of a degenerate induced module is called a *degenerate* irreducible module.

One of the most important problems of representation theory is to determine all degenerate representations. I will state here the result for $L = E(3|6)$ with $L_0 = \Pi_{j \geq 0} \mathfrak{a}_j$ (see Sect. 1.6.). The finite-dimensional irreducible L_0 -modules are actually $\mathfrak{a}_0 = \mathfrak{sl}_3 \oplus \mathfrak{sl}_2 \oplus \mathfrak{gl}_1$ -modules (with $\Pi_{j > 0} \mathfrak{a}_j$ acting trivially). We shall normalize the generator Y of $\mathfrak{gl}_1$ by the condition that its eigenvalue on $\mathfrak{a}_{-1}$ is $-1/3$. The finite-dimensional irreducible $\mathfrak{a}_0$ -modules are labeled by triples $(p, q; r; Y)$, where p, q (resp. r) are labels of the highest weight of an irreducible representation of $\mathfrak{sl}_3$ (resp. $\mathfrak{sl}_2$), so that $p, 0$ and $0, q$ label $S^p \mathbb{C}^3$ and $S^q \mathbb{C}^{3*}$ (resp. r labels $S^r \mathbb{C}^2$), and Y is the eigenvalue of the central element Y . Since irreducible $E(3|6)$ -modules are unique quotients of induced modules, they can be labeled by the above triples as well.

Theorem 4. [KR1]–[KR3] *The complete list of irreducible degenerate $E(3|6)$ -modules consists of four series: $(p, 0; r; -r + \frac{2}{3}p)$, $(p, 0; r; r + \frac{2}{3}p + 2)$, $(0, q; r; -r - \frac{2}{3}q - 2)$, $(0, q; r; r - \frac{2}{3}q)$.*

5.2. Remarkably, all four degenerate series occur as cokernels of the differential of a differential complex (M, ∇) constructed below (see [KR2] for details). I shall view $E(3|6)$ as a subalgebra of $E(5|10) = \Pi_{j \geq -2} \mathfrak{g}_j$ as in Sect. 1.6., expressed in terms of vector fields and differential forms in the indeterminates $x_1, x_2, x_3, z_+ = x_4, z_- = x_5$. Recall that $\mathfrak{g}_0$ is the algebra of divergenceless vector fields with linear coefficients. Let $Y = \frac{2}{3} \sum_i x_i \partial_i - \sum_\epsilon z_\epsilon \partial_\epsilon \in \mathfrak{g}_0$. Here and further $i = 1, 2, 3$, $\epsilon = +, -$, and $\partial_i = \partial/\partial x_i$, $\partial_\epsilon = \partial/\partial z_\epsilon$. Then $\mathfrak{a}_0$ is the centralizer of Y in $\mathfrak{g}_0$ and $\mathfrak{a}_{-1}$ is the span of all elements $d_i^\epsilon = dx_i \wedge dz_\epsilon$.

Consider the following four $\mathfrak{a}_0$ -modules (extended to L_0 -modules by trivial action of $\mathfrak{a}_j$ with $j > 0$):

$$V_I = \mathbb{C}[x_i, z_\epsilon], \quad V_{II} = \mathbb{C}[x_i, \partial_\epsilon]_{[2]}, \quad V_{III} = \mathbb{C}[\partial_i, z_\epsilon]_{[-2]}, \quad V_{IV} = \mathbb{C}[\partial_i, \partial_\epsilon],$$

where the subscript $[a]$ means that Y is shifted by the scalar a . For each $R = I - IV$ introduce a bigradation $V_R = \bigoplus_{m, n \in \mathbb{Z}} V_R^{(m, n)}$ by letting $\deg x_i = (1, 0)$, $\deg z_\epsilon = (0, 1)$, and let $M_R = \text{Ind}_{L_0}^L V_R = \bigoplus_{m, n \in \mathbb{Z}} M_R^{(m, n)}$. Then the non-zero $M_R^{(m, n)}$ are all the degenerate $E(3|6)$ -modules of the R^{th} series. We let

$$M = \left(\bigoplus_{(m, n) \neq (0, 0)} M_I^{(m, n)} \right) \oplus M_{II} \oplus M_{III} \oplus \left(\bigoplus_{(m, n) \neq (0, 0)} M_{IV}^{(m, n)} \right).$$

The differentials ∇_k introduced further are elements of $U(L) \otimes \text{End } V$ that act on $U(L) \otimes_{U(L_0)} V$ by the formula:

$$\left(\sum_j u_j \otimes A_j \right) (u \otimes v) = \sum_j u u_j \otimes A_j v.$$

Example. The dual to the ordinary formal de Rham complex is $\mathbb{C}[\partial/\partial x_1, \dots, \partial/\partial x_m] \otimes \Lambda(\partial/\partial \xi_1, \dots, \partial/\partial \xi_m)$ with the differential $d^* = \sum_j \partial/\partial x_j \otimes \xi_j$ and the $\mathbb{Z}$ -gradation defined by $\deg \partial/\partial x_i = 0$, $\deg \partial/\partial \xi_i = 1$. Rudakov's theorem [R] says that all irreducible degenerate W_m -modules occur as cokernels of d^* .

Turning now to ∇_k , we let $\Delta^\pm = \sum_i d_i^\pm \otimes \partial_i$, $\delta_i = d_i^+ \otimes \partial_+ + d_i^- \otimes \partial_-$. Then $\nabla_1 = \Delta^+(1 \otimes \partial_+) + \Delta^-(1 \otimes \partial_-)$ is a well-defined operator on all M_R such that $\nabla_1^2 = 0$. Furthermore there are differentials $\nabla_2 = \Delta^+ \Delta^-$, $\nabla_3 = \delta_1 \delta_2 \delta_3$, ∇_4 , ∇'_4 and ∇_6 (the explicit expressions of the last three can be found in [KR2]) that sew together these four complexes. These differentials are illustrated by Table M. The white nodes and black marks represent the induced modules of the R^{th} series. The plain arrows represent ∇_1 , the dotted arrows represent ∇_2 , the interrupted arrows represent ∇_3 and the bold arrows represent ∇'_4 , ∇''_4 and ∇_6 . The white nodes denote the places with zero homology. The black marks denote the places with non-zero homology, also computed in [KR2]. For example, at the star mark the homology is $\mathbb{C}$.

Similar results for $E(3|8)$ and (to a lesser extent) for $E(5|10)$ are given in [KR4].

5.3. The first hint that the Lie superalgebra $E(3|6)$ is somehow related to the Standard Model comes from the observation that its subalgebra $\mathfrak{a}_0$ is isomorphic to the complexified Lie algebra of the group of symmetries of the Standard Model. Table P below lists all $\mathfrak{a}_0$ -multiplets of fundamental particles of the Standard Model (see e.g. [O]): the upper part is comprised of three generations of quarks and the middle part of three generations of leptons (these are all fundamental fermions from which matter is built), and the lower part is comprised of the fundamental bosons (which mediate the strong and electro-weak interactions).

multiplets	charges	particles		
$(01, 1, \frac{1}{3})$	$\frac{2}{3}, -\frac{1}{3}$	$\begin{pmatrix} u_L \\ d_L \end{pmatrix}$	$\begin{pmatrix} c_L \\ s_L \end{pmatrix}$	$\begin{pmatrix} t_L \\ b_L \end{pmatrix}$
$(10, 1, -\frac{1}{3})$	$-\frac{2}{3}, \frac{1}{3}$	$\begin{pmatrix} \tilde{u}_R \\ \tilde{d}_R \end{pmatrix}$	$\begin{pmatrix} \tilde{c}_R \\ \tilde{s}_R \end{pmatrix}$	$\begin{pmatrix} \tilde{t}_R \\ \tilde{b}_R \end{pmatrix}$
$(10, 0, -\frac{4}{3})$	$-\frac{2}{3}$	$\tilde{u}_L$	$\tilde{c}_L$	$\tilde{t}_L$
$(01, 0, \frac{4}{3})$	$\frac{2}{3}$	u_R	c_R	t_R
$(01, 0, -\frac{2}{3})$	$-\frac{1}{3}$	d_R	s_R	b_R
$(10, 0, \frac{2}{3})$	$\frac{1}{3}$	$\tilde{d}_L$	$\tilde{s}_L$	$\tilde{b}_L$
$(00, 1, -1)$	$0, -1$	$\begin{pmatrix} \nu_L \\ e_L \end{pmatrix}$	$\begin{pmatrix} \nu_{\mu L} \\ \mu_L \end{pmatrix}$	$\begin{pmatrix} \nu_{\tau L} \\ \tau_L \end{pmatrix}$
$(00, 1, 1)$	$0, 1$	$\begin{pmatrix} \tilde{\nu}_R \\ \tilde{e}_R \end{pmatrix}$	$\begin{pmatrix} \tilde{\nu}_{\mu R} \\ \tilde{\mu}_R \end{pmatrix}$	$\begin{pmatrix} \tilde{\nu}_{\tau R} \\ \tilde{\tau}_R \end{pmatrix}$
$(00, 0, 2)$	1	$\tilde{e}_L$	$\tilde{\mu}_L$	$\tilde{\tau}_L$
$(00, 0, -2)$	-1	e_R	μ_R	τ_R
$(11, 0, 0)$	0	gluons		
$(00, 2, 0)$	$1, -1, 0$	W^+, W^-, Z (gauge bosons)		
$(00, 0, 0)$	0	γ (photon)		

Table P



Table M

It is easy to deduce from Theorem 4 that this list of multiplets (plus the multiplets $(11, 0, \pm 2)$) is characterized by the conditions:

- (i) $\mathfrak{a}_0$ -multiplet occurs in a degenerate irreducible $E(3|6)$ -module,
- (ii) when restricted to $sl_3 \subset \mathfrak{a}_0$, this multiplet contains only 1-dimensional, the two fundamental or the adjoint representation,
- (iii) $|Q| \leq 1$ for all particles of the multiplet, where the charge Q of a particle is given by the Gell-Mann-Nishijima formula: $Q = \frac{1}{2}(y + h)$, where y (resp. h) is the Y -eigenvalue (resp. $H = \text{diag}(1, -1) \in sl_2$ -eigenvalue).

How can we see the number of generations of quarks and leptons? For that order the sequence subcomplexes in Table M by $t = r - q + 3$ in sector IV (time), and replace in them the induced modules by their irreducible quotients. Then we find [KR2] (based on computer calculations by Joris Van der Jeugt) that a fundamental particle multiplet appears in the t^{th} sequence iff $t \geq 1$. Furthermore, for $1 \leq t \leq 7$ we get sequences with various particle contents, but for $t \geq 8$ the particle contents remains unchanged, and it is invariant under the CPT symmetry (though for $t \leq 7$ it is not). The explicit contents is exhibited in [KR2], 659–660.

Remarkably, precisely three generations of leptons occur in the stable region ($t \geq 8$), but the situation with quarks is more complicated: this model predicts a complete fourth generation of quarks and an incomplete fifth generation (with missing down type triplets).

In view of this discussion, it is natural to suggest that the algebra $su_3 + su_2 + u_1$ of internal symmetries of the Weinberg-Salam-Glashow Standard Model extends to $E(3|6)$. It is hoped that the representation theory of $E(3|6)$ will shed new light on various features of the Standard Model. I find it quite remarkable that the SU_5 Grand Unified Model of Georgi-Glashow combines the left multiplets of fundamental fermions in precisely the negative part of the consistent gradation of $E(5|10)$ (see Sec. 1.6). This is perhaps an indication of the possibility that the extension from su_5 to $E(5|10)$ algebra of internal symmetries may resolve the difficulties with the proton decay.

References

- [ALS] D. Alexseevski, D. Leites and I. Shchepochkina, Examples of simple Lie superalgebras of vector fields, *C.R.Acad.Bul.Sci.* **33** (1980), 1187–1190.
- [BPZ] A.A. Belavin, A.M. Polyakov and A.M. Zamolodchikov, Infinite conformal symmetry of critical fluctuations in two dimensions, *J.Stat.Phys.* **34** (1984), 763–774.
- [Be] M. Bershadsky, Conformal field theory via Hamiltonian reduction, *Comm. Math. Phys.* **139** (1991), 71–82.
- [BT] J. de Boer and T. Tjin, The relation between quantum W algebras and Lie algebras, *Comm. Math. Phys.* **160** (1994), 317–332.
- [B] R. Borcherds, Vertex algebras, Kac-Moody algebras, and the Monster, *Proc. Natl. Acad. Sci. USA*, **83** (1986), 3068–3071.
- [C] E. Cartan, Les groupes des transformations continués, infinis, simples, *Ann. Sci. Ecole Norm. Sup.* 26 (1909), 93–161.

- [CK1] S.-J. Cheng and V.G. Kac, A new $N = 6$ superconformal algebra, *Commun. Math. Phys.* **186** (1997), 219–231.
- [CK2] S.-J. Cheng and V.G. Kac, Generalized Spencer cohomology and filtered deformations of $\mathbb{Z}$ -graded Lie superalgebras, *Adv. Theor. Math. Phys.* **2** (1998), 1141–1182.
- [CK3] S.-J. Cheng and V.G. Kac, Structure of some $\mathbb{Z}$ -graded Lie superalgebras of vector fields, *Transformation Groups* **4** (1999), 219–272.
- [DK] A. D’Andrea and V.G. Kac, Structure theory of finite conformal algebras, *Selecta Mathematica* **4** (1998), 377–418.
- [DS] V.G. Drinfeld and V.V. Sokolov, Lie algebra and the KdV type equations, *Soviet J. Math* **30**, (1985), 1975–2036.
- [ET] T. Eguchi and A. Taormina, On the unitary representations of $N = 2$ and $N = 4$ superconformal algebras, *Phys Lett.* **B210** (1988), 125–132.
- [FK] D. Fattori and V.G. Kac, Classification of finite simple Lie conformal superalgebras, *J. Algebra* (2002).
- [FF1] B.L. Feigin and E.V. Frenkel, Representations of affine Kac-Moody algebras, bozination and resolutions, *Lett. Math. Phys.* **90** (1990), 307–317.
- [FF2] B.L. Feigin and E.V. Frenkel, Quantization of Drinfeld-Sokolov reduction, *Phys. Lett.* **B246** (1990), 75–81.
- [FST] B.L. Feigin, A.M. Semihkatov, V.A. Sirota and I. Yu Tipunin, Resolutions and characters of irreducible representations of the $N = 2$ superconformal algebra, hep-th/9805179.
- [FL] E.S. Fradkin and V. Ya. Linetsky, Classification of superconformal and quasisuperconformal algebras in two dimensions, *Phys. Lett.* **B291** (1992), 71–76.
- [FB] E. Frenkel and D. Ben-Zvi, “Vertex algebras and algebraic curves”, AMS monographs, **vol. 88**, 2001.
- [FKW] E. Frenkel, V. Kac and M. Wakimoto, Characters and fusion rules for W -algebras via quantized Drinfeld-Sokolov reduction, *Comm. Math. Phys.* **147** (1992), 295–328.
- [G1] V.W. Guillemin, A Jordan-Hölder decomposition for a certain class of infinite dimensional Lie algebras, *J. Diff. Geom.* **2** (1968), 313–345.
- [G2] V.W. Guillemin, Infinite-dimensional primitive Lie algebras, *J. Diff. Geom.* **4** (1970), 257–282.
- [GS] V.W. Guillemin and S. Sternberg, An algebraic model of transitive differential geometry, *Bull. Amer. Math. Soc.* **70** (1964), 16–47.
- [K1] V.G. Kac, Simple irreducible graded Lie algebras of finite growth, *Math. USSR-Izvestija* **2** (1968), 1271–1311.
- [K2] V.G. Kac, Lie superalgebras, *Adv. Math.* **26** (1977), 8–96.
- [K3] V.G. Kac, “Infinite-dimensional Lie algebras”, Third edition, Cambridge University Press, 1990.
- [K4] V.G. Kac, Modular invariance in mathematics and physics, in: “Mathematics into the 21st century”, *AMS Centennial Publ.*, 1992, 337–350.
- [K5] V.G. Kac, “Vertex algebras for beginners”, University Lecture Series, vol. 10 AMS, Providence, RI, 1996. Second edition 1998.

- [K6] V.G. Kac, Superconformal algebras and transitive group actions on quadrics, *Comm. Math. Phys.* **186** (1997), 233–252. Erratum, *Comm. Math. Phys.* **217** (2001), 697–698.
- [K7] V.G. Kac, Classification of infinite-dimensional simple linearly compact Lie superalgebras, *Adv. Math.* **139** (1998), 1–55.
- [K8] V.G. Kac, Classification of infinite-dimensional simple groups of supersymmetries and quantum field theory, “Visions in Mathematics toward 2000”, *GAFa*, Special volume (2000), 162–183.
- [KL] V.G. Kac and J.W. van de Leur, On classification of superconformal algebras, in: “Strings 88” (Eds S.J. Gates et al.), World Sci., 1989, 77–106.
- [KRW] V.G. Kac, S.-S. Roan and M. Wakimoto, Quantum reduction for affine Lie superalgebras, 2002 preprint.
- [KR1] V.G. Kac and A.N. Rudakov, Representations of the exceptional Lie superalgebra $E(3,6)$ I: Degeneracy conditions, *Transformation Groups* **7** (2002), 67–86.
- [KR2] V.G. Kac and A.N. Rudakov, Representations of the exceptional Lie superalgebra $E(3,6)$ II: Four series of degenerate modules, *Comm. Math. Phys.* **222** (2001), 611–661.
- [KR3] V.G. Kac and A.N. Rudakov, Representations of the exceptional Lie superalgebra $E(3,6)$ III: Classification of singular vectors.
- [KR4] V.G. Kac and A.N. Rudakov, Complexes of modules over exceptional Lie superalgebras $E(3,8)$ and $E(5,10)$, *IMRN* **19** (2002), 1007–1025.
- [KW1] V.G. Kac and M. Wakimoto, Modular invariant representations of infinite-dimensional Lie algebras and superalgebras, *Proc. Natl. Acad. Sci* **85** (1989), 4956–4960.
- [KW2] V.G. Kac and M. Wakimoto, Classification of modular invariant representations of affine superalgebras, in *Infinite-dimensional Lie algebras and groups*, Adv. Ser. Math. Phys. **vol. 7** World Scientific, 1989, 138–177.
- [KW3] V.G. Kac and M. Wakimoto, Integrable highest weight modules over affine superalgebras and number theory, *Progress in Math.* **Vol. 123** (1994), Birkhäuser Boston, 415–456.
- [KW4] V.G. Kac and M. Wakimoto, Integrable highest weight modules over affine superalgebras and Appell’s function, *Comm. Math. Phys* **215** (2001), 631–682.
- [KW5] V.G. Kac and M. Wakimoto, Quantum reduction and representation theory of superconformal algebras.
- [Kh] T. Khovanova, KdV superequation related to the Lie superalgebra of Neveu-Schwarz-2 string theory, *Teor. Mat. Phys.* **72** (1987), 899–904.
- [Ko] Yu. Kochetkoff, Déformations de superalgèbres de Buttin et quantification, *C.R. Acad. Sci. Paris* **299**, ser I, no. 14 (1984), 643–645.
- [KS] B. Kostant and S. Sternberg, Symplectic reduction, BRS cohomology and infinite-dimensional Clifford algebras, *Ann. Phys.* **176** (1987), 49–113.
- [L] S. Lie, Theorie der Transformations gruppen, *Math. Ann.* **16** (1880), 441–528.
- [M] K. Miki, The representation theory of the $SO(3)$ invariant superconformal

- algebra, *Int. J. Mod. Phys.* **5** (1990), 1293–1318.
- [O] L. Okun, “Physics of elementary particles”, Nauka, 1988 (in Russian).
- [P] A. Polishchuk, M.P. Appell’s function and vector bundles of rank 2 on elliptic curves, *Ramanujan J.* **5** (2001), 111–128.
- [RS] P. Ramond and J.H. Schwarz, Classification of dual model gauge algebras, *Phys.Lett.B.* **64** (1976), 75–77.
- [R] A.N. Rudakov, Irreducible representations of infinite-dimensional Lie algebras of Cartan type, *Math. USSR-Izvestija* **8** (1974), 836–866.
- [SV] J.A. Schouten and W. van der Kulk, “Pfaff problem and its generalizations”, Clarendon Press, 1949.
- [Sh1] I. Shchepochkina, New exceptional simple Lie superalgebras, *C.R. Bul. Sci.* **36**, no. 3 (1983), 313–314.
- [Sh2] I. Shchepochkina, Five exceptional simple Lie superalgebras of vector fields, *Funct. Anal. Appl.* **33** (1999), No. 3, 208–219.
- [Se] V. Serganova, Characters of irreducible representations of simple Lie superalgebras, *Doc. Math.*, Extra Volume ICM 1998 II, 583–593.
- [Wa1] M. Wakimoto, Representation theory of affine superalgebras at the critical level, *Doc. Math.* Extra Volume ICM 1998 II, 605–614.
- [Wa2] M. Wakimoto, “Lectures on infinite-dimensional Lie algebras”, *World Sci.*, 2001.
- [W] B. Yu Weisfeiler, Infinite-dimensional filtered Lie algebras and their connection with graded Lie algebras, *Funct. Anal. Appl.* **2** (1968), 88–89.
- [Z] A.B. Zamolodchikov, Infinite additional symmetries in 2-dimensional conformal quantum field theory *Teor. Mat. Phys.* **65** (1985), 1205–1213.